

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 49/00, 49/04	A1	(11) International Publication Number: WO 95/21631 (43) International Publication Date: 17 August 1995 (17.08.95)
(21) International Application Number: PCT/GB95/00316 (22) International Filing Date: 15 February 1995 (15.02.95) (30) Priority Data: 9402867.7 15 February 1994 (15.02.94) GB (71) Applicant (for all designated States except US): NYCOMED IMAGING AS [NO/NO]; Nycoveien 2, P.O. Box 4220 Torshov, N-0401 Oslo 4 (NO). (71) Applicants (for GB only): HOLMES, Michael, John [GB/GB]; Imperial House, 15-19 Kingsway, London WC2B 6UZ (GB). HOLMES, Michael, John [GB/GB]; Imperial House, 15-19 Kingsway, London WC2B 6UZ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): DUGSTAD, Harald [NO/NO]; Jupiterveien 13, N-0489 Oslo (NO). FOSS, Per, Antonius [NO/NO]; Konventveien 3B, N-0377 Oslo (NO). KLAVENESS, Jo [NO/NO]; Midtåsen 5, N-1166 Oslo (NO). RONGVED, Pål [NO/NO]; Hovdens vei 11, N-1450 Nesoddtangen (NO). SKURTVEIT, Roald [NO/NO]; Schweigaards gate 48B, N-0191 Oslo (NO). SOLBERG, Jan [NO/NO]; Syd-Fossum 43, N-1343 Eiksmarka (NO).	AUKRUST, Inger, Reidun, Fjeldskaar [NO/NO]; Stokkevn 5, N-3400 Lier (NO). (74) Agent: MARSDEN, John, Christopher; Frank B. Dehn & Co., Imperial House, 15-19 Kingsway, London WC2B 6UZ (GB). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UG, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published With international search report.	
(54) Title: IMPROVEMENTS IN OR RELATING TO CONTRAST AGENTS		
(57) Abstract Microparticulate contrast agents comprising gas or a gas precursor encapsulated by a non-polymeric and non-polymerisable wall-forming material are readily characterisable materials exhibiting surprising structural integrity and stability.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

"Improvements in or relating to contrast agents"

5 This invention relates to novel contrast agents,
more particularly to new gas-containing and gas-
generating contrast agents of use in diagnostic imaging,
and to methods for their preparation and use.

10 It is well known that ultrasonic imaging comprises
a potentially valuable diagnostic tool, for example in
studies of the vascular system, particularly in
cardiography, and of tissue microvasculature. A variety
of contrast agents has been proposed to enhance the
acoustic images so obtained, including suspensions of
solid particles, emulsified liquid droplets, gas bubbles
15 and encapsulated gases or liquids. It is generally
accepted that low density contrast agents which are
easily compressible are particularly efficient in terms
of the acoustic backscatter they generate, and
considerable interest has therefore been shown in the
20 preparation of gas-containing and gas-generating
systems.

Gas-containing contrast media are also known to be
effective in magnetic resonance (MR) imaging, e.g. as
susceptibility contrast agents which will act to reduce
25 MR signal intensity. Oxygen-containing contrast media
also represent potentially useful paramagnetic MR
contrast agents.

Furthermore, in the field of X-ray imaging it has
been observed that gases such as carbon dioxide may be
30 used as negative oral contrast agents.

Initial studies involving free gas bubbles
generated in vivo by intracardiac injection of
physiologically acceptable substances have demonstrated
the potential efficiency of such bubbles as contrast
35 agents in echocardiography; such techniques are severely

- 2 -

limited in practice, however, by the short lifetime of the free bubbles. Interest has accordingly been shown in methods of stabilising gas bubbles for echocardiography and other ultrasonic studies, for example using emulsifiers, oils, thickeners or sugars, or by entraining or encapsulating the gas or a precursor therefore in a variety of systems, e.g. as porous gas-containing microparticles or as encapsulated gas microbubbles.

Previous proposals relating to microbubble-containing ultrasound contrast agents have invariably required the use of a polymeric encapsulating coating for the microbubbles. Thus, for example, WO-A-8002365, which in principle suggests use of microbubbles having a coalescence resistant encapsulating membrane comprising non-toxic and non-antigenic organic molecules, in practice discloses only the use of gelatin as the encapsulating material. It has been found that microbubbles so encapsulated do not exhibit adequate stability at the dimensions preferred for use in echocardiography (1-10 μm) in view of the extreme thinness of the encapsulating coating.

US-A-4774958 discloses the use of microbubble dispersions stabilised by encapsulation in denatured protein, e.g. human serum albumin. Such systems permit the production of microbubble systems having a size of e.g. 2-5 μm but still do not permit efficient visualisation of the left heart and myocardium. The use of such protein-derived agents may also create problems with regard to potential allergenic reactions.

EP-A-0327490 and WO-A-8906978 disclose, inter alia, ultrasonic contrast agents comprising microparticulate amylose or a synthetic biodegradable polymer containing a gas or volatile fluid (i.e. having a boiling point below 60°C) in free or bonded form. Representative synthetic biodegradable polymers include polyesters of hydroxy carbonic acids, polyalkyl cyanoacrylates,

- 3 -

polyamino acids, polyamides, polyacrylated saccharides and polyorthoesters.

Similar biodegradable microparticulate polymers, based on polymerised aldehydes, are described in EP-A-0441468, while systems based on microparticulate poly (amino acid) - poly (cyclic imide) derivatives are described in EP-A-0458079, US-A-5137928, US-A-5190982, US-A-5205287 and US-A-5229469.


EP-A-0458745 discloses air or gas-filled microballoons in which the encapsulating material is a deformable and resilient interfacially deposited polymer which is preferably biodegradable, examples including polysaccharides, polyamino acids, polylactides, polyglycolides, lactide/lactone copolymers, polypeptides, proteins, polyorthoesters, polydioxanone, poly- β -aminoketones, polyphosphazenes, polyanhydrides and poly (alkyl cyanoacrylates). The microballoons are normally prepared by emulsion techniques leading to deposition of the polymer around droplets of a volatile liquid which is subsequently evaporated. Such techniques generally involve the use of surfactants, for example lecithins, fatty acids or esters thereof with polyoxyalkylene compounds such as polyoxyethylene glycol or polyoxypropylene glycol, in order to stabilise the emulsion.

It is generally acknowledged that polymer-based contrast agents should desirably be biodegradable in order to facilitate their ultimate elimination from or absorption by the test subject. In many instances it has therefore been proposed to use polymers such as polyesters, polyanhydrides, polycarbonates, polyamides and polyurethanes which are biodegradable as a result of the susceptibility of ester, amide or urethane groups therein to enzymic hydrolysis in vivo.

In WO-A-9317718 there are described polymer-based contrast agents which are designed to exhibit high and controllable levels of biodegradability in vivo by

- 4 -

virtue of the presence in the polymer of methylene diester units of formula (I)



5 (where R^1 and R^2 each represent a hydrogen atom or a carbon-attached monovalent organic group or R^1 and R^2 together form a carbon-attached divalent organic group and m and n , which may be the same or different, are
10 each zero or 1). Such units are particularly rapidly degraded by common esterase enzymes but are relatively stable in the absence of enzymes.

In all the above-described encapsulated microbubble contrast agents the material encapsulating the
15 microbubbles consists essentially of polymer, although minor quantities of other materials may be present. Thus, for example, EP-A-0458745 suggests that additives such as fats, waxes, high molecular weight hydrocarbons, phospholipids and plasticisers may be incorporated into
20 the polymer wall, e.g. in amounts of up to 20% by weight. Clearly, however, it has hitherto been thought necessary to employ polymeric encapsulating material, e.g. in order to achieve sufficient structural integrity so as to impart adequate stability to the contrast
25 agent. An isolated exception is WO-A-9401140, which discloses micro-gas bubble-containing echographic contrast agents prepared by lyophilising aqueous emulsions containing a lipid-soluble or water-insoluble builder such as cholesterol; these contrast agents are,
30 however, also required to contain a substantial proportion of an apolar liquid such as petroleum ether.

We have now most surprisingly found that effective contrast agents comprising encapsulated microbubbles may be prepared using a wide range of non-polymeric
35 wall-forming materials to encapsulate the gas or a precursor therefor. It will be appreciated that such contrast agents may exhibit significant advantages over

- 5 -

polymer-based contrast agents, in particular that they may be easier and more economical to prepare and easier to characterise; they may also be more readily eliminable from the bodies of subjects to whom they are administered, for example by virtue of the smaller size and/or enhanced biodegradability of the non-polymeric molecules. The selection of materials which are endogeneous or are biodegradable to endogenous substances may also be advantageous.

10 Thus according to one aspect of the present invention there is provided a microparticulate contrast agent comprising gas or a gas precursor encapsulated by a non-polymeric and non-polymerisable wall-forming material.

15 The term "non-polymeric" as used herein denotes that the wall-forming materials do not contain multiple repeating units joined head-to-tail, as in polymers, and are not obtained by polymerisation techniques. The wall-forming materials will thus most commonly comprise
20 well-defined and characterisable molecules, e.g. as evidenced by precise melting points, single chromatographic mobilities etc., and by monodisperse molecular weights (i.e. having a polydispersity index of 1.0, where this is defined as the ratio of weight
25 average molecular weight to number average molecular weight), although it will be appreciated that the invention also embraces, for example, contrast agents comprising clearly constituted mixtures of such substances, including naturally occurring mixtures.
30 These properties may be contrasted with those of materials obtained by polymerisation techniques, where products will typically comprise variable mixtures of molecules of different chain length and so will be characterised by a melting point range, a plurality of
35 chromatographic mobilities and a molecular weight range (e.g. as evidenced by a polydispersity index higher than 1.0).

- 6 -

The term "non-polymerisable" indicates that the encapsulating material will not polymerise under, for example, conditions such as those used during preparation and storage of the contrast agents of the invention.

The encapsulating material may conveniently have a low molecular weight, for example in the range 100-6000. It will be appreciated that the molecular weight should be such that the melting point and solubility properties of the material permit microparticle formation at an appropriate temperature and under appropriate processing conditions, e.g. as described hereinafter.

It will be appreciated that in order to exhibit the necessary wall-forming properties the encapsulating material should be solid or semi-solid at normal storage and handling temperatures; the material therefore advantageously has a melting point of at least 40°C; in many cases such materials will exhibit film-forming properties, which may be enhanced by virtue of the material having a somewhat amphiphilic character as a result of the presence of hydrophilic and lipophilic regions within the molecules; the hydrophilic character may for example derive from specifically hydrophilic groups such as carboxy, keto, hydroxy or amino and/or from the presence of groups enabling participation in hydrogen bonding, for example the carbonyl moieties of esterified carboxy groups.

In order to facilitate preparation of the contrast agents of the invention, e.g. by procedures such as those described hereinafter, the encapsulating material desirably has greater oil solubility than aqueous solubility.

In addition, the encapsulating material and any degradation products thereof which may be generated in vivo must be physiologically compatible and should either be endogenous or readily eliminable, optionally after breakdown into smaller molecules, e.g. as a result

- 7 -

of hydrolytic, enzymatic or other metabolic reactions, for example involving labile linkages such as ester, amide or urethane groups.

The contrast agents of the invention may comprise a wide range of wall-forming materials which fulfil the above-described requirements, including, for example, fatty acids, e.g. lipophilic saturated or unsaturated aliphatic carboxylic acids containing 10-50 carbon atoms, e.g. 10-30 carbon atoms, such as palmitic, stearic or behenic acid, and esters thereof, e.g. alkyl esters, for example polyhydroxyalkyl esters such as pentaerythritol, ethylene glycol or glyceryl esters; fatty alcohols and amines, e.g. lipophilic saturated or unsaturated aliphatic alcohols and amines containing 10-50 carbon atoms. e.g. 10-30 carbon atoms, and esters or amides thereof, e.g. with mono-, di- or tri-carboxylic acids, for example optionally hydroxylated lower alkanolic acids such as acetic, adipic and citric acid; lipophilic aldehydes and ketones; lipophilic derivatives of sugars, e.g. containing lipophilic ether or, more preferably, ester groups; cholic acids and derivatives (e.g. esters) thereof; cholesterol and derivatives thereof; aliphatic and aromatic hydrocarbons such as mineral waxes; hydrophobically modified hydrophilic compounds, e.g. hydrophilic compounds such as X-ray contrast agents modified to contain one or more lipophilic groups (e.g. hydrocarbyl groups such as aliphatic chains) so as to render the molecule hydrophobic; and other biocompatible fat-soluble materials, e.g. antioxidants such as tocopherols or thiocctic acid and derivatives thereof.

Contrast agents comprising wall-forming materials containing one or more methylene diester units (e.g. of formula (I) above) per molecule are useful embodiments of the invention by virtue of the ease with which they may be cleaved in vivo to form smaller and more readily eliminable molecules. Such units may also impart a

- 8 -

desirable amphiphilic character to the wall-forming material.

Any biocompatible gas may be employed in the contrast agents of the invention, the term "gas" as used herein embracing any substance in gaseous form at 37°C. Representative gases include air, nitrogen, oxygen, hydrogen, nitrous oxide, carbon dioxide, helium, argon and low molecular weight hydrocarbons such as methane and acetylene. Low molecular weight fluorinated compounds such as sulphur hexafluoride, disulphur decafluoride, carbon tetrafluoride and perfluoroalkanes such as perfluoropropane, perfluorobutane and perfluoropentane may be of particular interest. The gas may be free within the microbubbles or may be trapped or entrained within a containing substance.

Contrast agents according to the invention may be used in a variety of diagnostic imaging techniques, including ultrasound, MR and X-ray imaging. Their use in diagnostic ultrasound imaging and in MR imaging, e.g. as susceptibility contrast agents, constitute preferred features of the invention.

For ultrasound applications such as echocardiography, in order to permit free passage through the pulmonary system and to achieve resonance with the preferred imaging frequency of about 0.1-15 MHz, it may be convenient to employ microparticles having an average size of 0.1-10 μ m, e.g. 1-7 μ m. Substantially larger particles or bubbles, e.g. with average sizes of up to 500 μ m, may however be useful in other applications, for example gastrointestinal imaging or investigations of the uterus or Fallopian tubes.

The contrast agents of the invention may be formulated in any manner appropriate to the intended method of administration, e.g. as suspensions in injectable media such as sterile water for injection. Such formulations may if desired contain biocompatible additives, e.g. antioxidants such as tocopherols or

- 9 -

thioctic acid.

The contrast agents of the invention may be prepared by any convenient method, for example by techniques analogous to those described in WO-A-9317718.

5 Representative techniques for the preparation of materials encapsulated by a wall or membrane are also described in literature such as "Microencapsulation and Related Drug Processes" by P.D. Deasy, Marcel Dekker Inc., New York (1984).

10 Thus, for example, contrast agents according to the invention may be prepared by emulsion techniques analogous to those in the polymer art. Typically such processes may involve (i) generating an emulsion comprising hydrophilic and hydrophobic phases wherein
15 the wall-forming material is preferentially solubilised in the dispersed phase or is distributed about the interfaces between the phases, and (ii) isolating the desired contrast agent from the emulsion. The emulsion will preferably also comprise one or more emulsifiers
20 solubilised in either or both phases. Single or multiple emulsions may be generated; representative multiple emulsion techniques are described in WO-A-9317718.

The wall-forming material may be selected to have a
25 lyophilicity appropriate to a particular form of emulsion processing; thus, for example, it may be advantageous to select an oil-soluble wall-forming material for processing using an oil-in-water emulsion. Oil-in-water emulsions may also be used to process wall-
30 forming materials having a degree of water-solubility where the material exhibits attractive interactions sufficient to slow the kinetics of dissolution so as to permit microparticle formation in the presence of water; a similar approach may be taken using a water-in-oil
35 emulsion and a wall-forming material having a degree of oil-solubility.

Emulsions may be prepared by, for example,

- 10 -

conventional techniques such as agitation, sonication, stirring (preferably high speed stirring) or other forms of mixing (e.g. high shear mixing), membrane emulsification, high-voltage emulsification or high pressure homogenisation. The wall-forming material may advantageously be predissolved in what is to be the dispersed phase. As noted above, one or more emulsifiers may advantageously be solubilised in either or both phases of the emulsion; preferably at least one emulsifier is solubilised in the continuous phase.

It will be appreciated that factors such as stirring speed will influence the size of the encapsulated microbubbles ultimately produced; thus, for example, faster stirring will tend to yield smaller microbubbles. The amount of emulsifier employed may vary widely, but typically may equal or exceed the amount of wall-forming material on a weight basis. Additives such as emulsification assistants, e.g. viscosity enhancers such as proteins, carbohydrates, polysaccharides or other hydrophilic polymers, may if desired also be employed.

In an alternative procedure a solution of wall-forming material in an appropriate aprotic organic solvent (e.g. a sulfoxide such as dimethyl sulfoxide, a cyclic ether such as tetrahydrofuran or an N,N-disubstituted amide such as dimethylformamide) may be mixed with an aqueous phase (e.g. using a high speed stirrer) so as to precipitate wall-forming material, which may be collected and lyophilised to yield the desired contrast agent. The aqueous phase may advantageously contain a polymeric material such as polyvinyl alcohol or a poloxamer (e.g. a Pluronic). Such techniques are described in the above-mentioned EP-A-0458079.

A further process comprises injecting a solution of the wall-forming material in an appropriate aprotic solvent into liquid nitrogen; the solution may, if

- 11 -

desired, also contain an additive such as hydroxypropylcellulose. Alternatively the wall-forming material may be dissolved in an appropriate solvent or dispersed in, for example, an oil-in-water, water-in-oil or multiple emulsion, and the solution or emulsion spray
5 dried, e.g. as described in EP-A-0514970.

Coacervation techniques, e.g. as are known in the art, may also be employed in preparing contrast agents according to the invention.

10 In principle any emulsifier may be used in the preparation of contrast agents according to the invention. We have found that it may, however, be advantageous to select a polymeric emulsifier, since the stability of the contrast agents so obtained may thereby
15 be enhanced, e.g. as evidenced by long-lasting retention of gas content coupled with a lower or minimal tendency of the microparticles to aggregate. Examples of such polymeric emulsifiers include polyvinyl alcohol, proteins (e.g. gelatin, albumins such as human or
20 porcine serum albumin, and casein salts such as sodium caseinate), polysaccharides (e.g. modified chitosan, modified starches including lipophilised starch and soluble reduced amylose, heparin, and gums such as gum arabic), block copolymers consisting of alternating
25 hydrophilic and hydrophobic blocks (e.g. polyhydroxystearate - polyethylene oxide block copolymers such as B246 (117753-68-1), polyoxyethylene - polyoxypropylene block copolymers, including poloxamers
such as tetronics and Pluronic F38, F68, F77, F88, F127,
30 L44, L64, P84 or P123, or perfluorinated derivatives thereof), polyoxyethylated derivatives of partial fatty acid esters of hexahydric alcohols such as sorbitol (e.g. Tween-type surfactants such as Tween 40, 60 or 80), polyvinylpyrrolidones (e.g. Kollidon), fatty
35 alcohol ethers of polyoxyethylene alcohols (e.g. Brij-type surfactants such as Brij 52 or 99), polyethylene glycol esters of fatty acids (e.g.

- 12 -

Cremaphor-type surfactants such as Cremaphor RH40), polyethylene glycol - sorbitan - beeswax surfactants such as Atlas G-1702 (PEG-6 sorbitan beeswax) and Atlas G-1726 (PEG-20 sorbitan beeswax), and perfluorinated derivatives of any of the above polyethylene glycol -
5 containing compounds.

While we do not wish to be bound by theoretical considerations, it may be that such polymeric emulsifiers are particularly compatible with the
10 wall-forming material and may, for example, be present as external coatings on the encapsulating material in contrast agents according to the invention. As such they may, for example, enhance the subsequent dispersibility and stability of the contrast agents,
15 e.g. inhibiting aggregative tendencies through electrostatic or other interactions. It will be appreciated that in such contrast agents the role of the emulsifier is as an emulsification aid and/or stabiliser and that it does not act as a wall-forming material.

20 It will be appreciated that the water-immiscible solvent used in emulsification procedures to prepare contrast agents according to the invention should be liquid at the temperature used for the emulsification process; it may advantageously be selected to have a
25 vapour pressure which facilitates its removal at a later stage. Representative solvents include aliphatic, cycloaliphatic and araliphatic hydrocarbons, e.g. containing up to 16 carbon atoms, such as n-octane, cyclooctane, a dimethylcyclohexane, ethylcyclohexane, a
30 methylheptane, an ethylhexane, toluene, xylene, naphthalene or a terpene, terpenoid or isoprenoid such as camphene or limonene; haloalkanes such as Freons, methylene chloride, chloroform, carbon tetrachloride or methyl bromide; esters such as ethyl or propyl acetate,
35 butyl formate and propyl or isopropyl butyrate or isobutyrate; and appropriate ethers and other substances which are liquid and have appropriate dissolving

- 13 -

properties at the temperature used for the emulsification process.

Where in the preparation of contrast agents according to the invention it is desired to deposit wall forming material at the interface between the oil and water phases of an emulsion, such deposition may be induced by, for example, modifying parameters such as temperature, pH, solvent properties or solute concentrations. Thus, for example, it is possible to induce the wall-forming material to precipitate by reducing the temperature of the emulsion, by pH change, by adding a non-solvent for the wall-forming material, or by at least partial removal of at least the organic solvent (e.g. by evaporation or lyophilisation, preferably under an atmosphere of the gas which is desired to be incorporated, for example as described in EP-A-0458745).

Where it is desired to prepare a gas precursor-containing contrast agent the precursor may, for example, conveniently be dissolved in the water-immiscible organic solvent prior to emulsification. The gas precursor may, for example, be a compound which reacts to produce gas following administration to a subject, e.g. as a result of decomposition induced thermally or by pH change or as a result of enzymatic degradation. Thus, for example, non-toxic organic carbonates and bicarbonates, e.g. arginine carbonate and compounds of formula $RO.CO.OM$ where R is an organic group and M represents a physiologically acceptable cation, will generate carbon dioxide in the conditions of pH prevailing in the bloodstream, as will compounds such as aminomalonates.

In general, the contrast agent may be isolated by any convenient method, for example by solvent removal using techniques such as evaporation, lyophilisation or spray drying. Such procedures may if desired be carried out under an atmosphere of the gas which is to be

- 14 -

incorporated in the contrast agent, if desired at reduced pressure.

- Contrast agents so obtained may be stored and transported in dry form, in which condition they may be
- 5 stable indefinitely, being mixed with an appropriate liquid carrier (e.g. sterile water for injection, physiological saline or phosphate buffer) prior to administration. In this way the concentration of the injected or otherwise administered contrast agent may be
- 10 varied at will depending on the precise nature of the application. The contrast agents may also be stored as suspensions in such carriers, especially where the porosity of the encapsulating membrane is comparatively low.
- 15 The following non-limitative Examples serve to illustrate the invention.

- 15 -

EXAMPLE 1 - SYNTHESIS OF WALL-FORMING MATERIALS AND EMULSIFIERS

5

a) Methylene bis(16-hexadecanoyloxyhexadecanoate)

A mixture of methylene bis(16-hydroxyhexadecanoate) (1.0 mmol) and palmitoyl chloride (7.0 mmol) was refluxed in diethyl ether (10 ml) for 24 hours before the solvent was evaporated. The residual material was purified on a silica gel flash column eluting first with hexane/chloroform (2:1) and then with chloroform. Yield 88%, white solid. ^1H NMR (CDCl_3): δ 0.88 (6 H, t, J 6.7 Hz), 1.2-1.4 (92 H, m), 1.5-1.7 (12 H, m), 2.28 (4 H, t, J 7.5 Hz), 2.35 (4 H, t, J 7.6 Hz), 4.05 (4 H, t, J 6.7 Hz), 5.75 (2 H, s). ^{13}C NMR (CDCl_3): δ 14.11, 22.70, 24.64, 25.06, 25.97, 28.70, 29.04, 29.19, 29.25, 29.29, 29.38, 29.47, 29.50, 29.56, 29.62, 29.67, 29.70, 29.71, 31.94, 34.00, 34.43, 64.39, 79.06, 172.59, 173.97.

b) Methylene (16-hexadecanoyloxyhexadecanoate) (16'-hydroxyhexadecanoate)

A mixture of methylene bis(16-hydroxyhexadecanoate) (1.0 mmol) and palmitoyl chloride (1.0 mmol) was refluxed in diethyl ether (10 ml) for 24 hours before the solvent was evaporated. The residual material was purified by repeated filtration through silica gel using methylene chloride/acetonitrile (95:5) for elution. Yield 38%, white solid. ^1H NMR (CDCl_3): δ 0.88 (3 H, t, J 6.7 Hz), 1.2-1.4 (69 H, m), 1.5-1.7 (10 H, m), 2.29 (2 H, t, J 7.5 Hz), 2.35 (4 H, t, J 7.6 Hz), 3.64 (2 H, t, J 6.6 Hz), 4.05 (2 H, t, J 6.7 Hz), 5.75 (2 H, s). ^{13}C NMR (CDCl_3): δ 14.12, 22.71, 24.64, 25.06, 25.77, 25.97, 28.70, 29.04, 29.19, 29.24, 29.25, 29.30, 29.38, 29.46, 29.50, 29.56, 29.60, 29.61, 29.63, 29.64, 29.66, 29.67,

- 16 -

29.70, 29.71, 31.95, 32.85, 34.01, 34.44, 63.09, 64.41, 79.06, 172.51, 174.01.

- 5 c) Product from reaction of methylene (16-hexadecanoyloxyhexadecanoate) (16'-hydroxyhexadecanoate) with adipoyl chloride

A solution of methylene (16-hexadecanoyloxyhexadecanoate) (16'-hydroxyhexadecanoate) (1.0 mmol) and
10 adipoyl chloride (0.5 mmol) was refluxed in xylene/trichloroethylene (4:1, 20 ml) at 60°C under reduced pressure (150 mBar) for 16 hours before the solvent was evaporated. The crude product was purified by flash chromatography on silica gel using methylene chloride/
15 acetonitrile (97:3) for elution. Yield 74%, white solid. ¹H NMR (CDCl₃): δ 0.88 (6 H, t, J 6.7 Hz), 1.2-1.4 (136 H, m), 1.5-1.7 (24 H, m), 2.28 (4 H, t, J 7.5 Hz), 2.35 (12 H, t, J 7.6 Hz), 4.05 (8 H, t, J 6.7 Hz), 5.74 (4 H, s). ¹³C NMR (CDCl₃): δ 14.12, 22.71, 24.48,
20 24.64, 25.06, 25.95, 25.97, 28.68, 28.70, 29.04, 29.19, 29.25, 29.30, 29.38, 29.47, 29.50, 29.56, 29.62, 29.67, 29.70, 29.72, 31.95, 33.98, 34.00, 34.43, 64.40, 64.56, 79.06, 172.49, 173.41, 173.98.

- 25 d) Methylene bis(16-hexadecanoyloxymethoxycarbonyloxy-hexadecanoate)

- 30 i) Methylene bis(16-chloromethoxycarbonyloxy-hexadecanoate)

Chloromethyl chloroformate (0.93 g, 7.2 mmol) was added to an ice-cooled solution of methylene bis(16-hydroxyhexadecanoate) (2.0 g, 3.6 mmol) in methylene
35 chloride (80 ml). Pyridine (0.57 g, 7.2 mmol) was added and the reaction mixture was stirred for 15 minutes at 0°C and 4 hours at room temperature. The reaction

- 17 -

mixture was washed with hydrochloric acid (1M, 50ml) saturated aqueous sodium bicarbonate (50 ml), water (50 ml) and dried (MgSO₄). The solvent was evaporated and the residue was purified by flash chromatography (silica/methylene chloride). Yield: 84% ¹H NMR (200 MHz, CDCl₃) : δ 1.25 (m, 44H), 1.72 (m, 8H), 2.31 (t, 4H), 4.20 (t, 4H), 5.72 (m, 6H).

ii) Methylene bis(16-hexadecanoyloxymethoxycarbonyl-oxyhexadecanoate)

Potassium t-butoxide (0.784 g, 7.0 mmol) was added to a solution of palmitic acid (1.80 g, 7.0 mmol) in N,N-dimethylformamide (100 ml), Methylene bis(16-chloromethoxycarbonyloxyl-oxyhexadecanoate) (2.6 g, 3.5 mmol) in N,N-dimethylformamide (10 ml) was added to the resulting suspension, followed by 18-crown-6 (0.1g). The reaction mixture was stirred at room temperature for 2 days. The reaction mixture was filtered and the solvent was removed under reduced pressure. The residue was purified by flash chromatography (silica/methylene chloride). Yield: 19%. ¹H NMR (300MHz, CDCl₃): δ 0.86 (m, 6H), 1.23 (m, 12H), 1.62 (m, 92H), 2.33 (m, 8H), 4.16 (m, 4H), 5.73 (m, 6H).

e) 1-(Octadecyloxycarbonyloxy)ethyl 5-acetamido-3-(N-methylacetamido)-2,4,6-triiodobenzenecarboxylate

i) 1-Chloroethyl octadecyl carbonate

30

1-Chloroethyl chloroformate (7.15g, 50mmol) and 1-octadecanol (13.53g, 50 mmol) were suspended in chloroform (100ml) at 0° C. Pyridine (3.96g, 50mmol) was added dropwise during 20 minutes, maintaining the temperature below 10°C. After stirring at room temperature for 24 hours the reaction mixture was washed

- 18 -

four times with 1N hydrochloric acid, once with a saturated sodium bicarbonate solution and finally twice with water. The organic solution was dried with magnesium sulphate and the solvent was removed at reduced pressure. Yield: 19g.

ii) 1-(Octadecyloxy carbonyloxy)ethyl 5-acetamido-3-(N-methylacetamido)-2,4,6-triiodobenzenecarboxylate

10 1-Chloroethyl octadecyl carbonate (7.54g, 20mmol) was added at room temperature to a solution of potassium 5-acetamido-3-(N-methylacetamido)-2,4,6-triiodobenzenecarboxylate (15.99g, 24mmol) and potassium iodide (0.33g, 2mmol) in dry N,N-dimethylformamide.

15 After stirring at 50°C for 18 hours the solvent was removed at reduced pressure. The residue was suspended in chloroform (200ml) and washed three times with a saturated sodium bicarbonate solution and finally twice with water. After drying with magnesium sulphate the

20 solvent was removed at reduced pressure. Yield: 18.6g. The product was further purified by flash chromatography (Silicagel 60, chloroform/acetonitrile 85:15). ¹H NMR (DMSO-d₆): δ 0.852 (CH₂CH₃), 1.236 (CH₂), 1.639 (CHCH₃), 1.657 (COOCH₂CH₂), 1.670 (N(CH₃)COCH₃), 2.051 (NCOCH₃), 2.965 (NCH₃), 4.154 (COOCH₂), 6.971 (CHCH₃), 10.103 (NH).

f) Methylene bis[3-(2,3-dihexadecanoyloxy-propoxycarbonyl)propionate]

30

i) 3-(2,3-Dihexadecanoyloxypropoxycarbonyl)propionic acid

Dipalmitin (1,2-dipalmitoylglycerol), (3.00 g, 5.27 mmol), succinic anhydride (1.00 g, 10.0 mmol) and N,N-dimethyl-4-aminopyridine (50 mg, 0.41 mmol) were dissolved in tetrahydrofuran (30 ml) and N,N-dimethylformamide (10

- 19 -

ml) under a nitrogen atmosphere. Triethylamine (2.5 ml) was subsequently added and the resulting solution was stirred for two days at ambient temperature. The solvents were evaporated at reduced pressure and the white solid residue was redissolved in chloroform (150 ml). The solution was washed with 1% aqueous hydrochloric acid (10 ml) and with brine (2x50ml), dried with magnesium sulphate and finally concentrated, yielding 5.80 g of a white solid which was used without further purification in the subsequent reaction.

¹H NMR (300 MHz, CDCl₃): δ 0.85-0.90 (m, 6 H), 1.20-1.35 (m, 48 H), 1.55-1.65 (m, 4 H), 2.30 (dd, J₁=7.6, J₂=2.5 Hz, 2 H), 2.32 (dd, J₁=7.4, J₂=2.3 Hz, 2 H), 2.62-2.70 (m, 4 H), 4.14 (dd, J₁=12.1, J₂=5.9 Hz, 1 H), 4.18 (dd, J₁=12.1, J₂=5.9 Hz, 1 H), 4.29 (dd, J₁=9.0, J₂=4.4 Hz, 1 H), 4.33 (dd, J₁=9.0, J₂=4.4 Hz, 1H), 5.23-5.30 (m, 1H).

¹³C NMR (300 MHz, CDCl₃): δ 14.121, 22.720, 24.890, 24.917, 28.716, 28.756, 29.110, 29.156, 29.310, 29.397, 29.517, 29.530, 29.697, 29.731, 31.961, 34.077, 34.211, 62.058, 62.673, 68.802, 171.620, 172.956, 173.349, 177.162.

ii) Methylene bis[3-(2,3-dihexadecanoyloxypropoxy-carbonyl)propionate]

1,8-Diazabicyclo[5.4.0]undec-7-ene (633 mg, 4.158 mmol) was added to a solution of 3-(2,3-dihexadecanoyloxypropoxycarbonyl)propionic acid from (a) above (2.65g, 3.96 mmol) in dry methylene chloride and N,N-dimethylformamide (40 ml, 1:1) under a nitrogen atmosphere. After 15 minutes stirring of the solution, diiodomethane (530 mg, 1.980 mmol) was added. The resulting solution was stirred at ambient temperature for four days, the solvents were evaporated at reduced pressure and the solid residue was redissolved in methylene chloride (200 ml). The solution was washed with 1% aqueous hydrochloric acid (1x25ml) and brine

- 20 -

(3x50ml), dried with magnesium sulphate and concentrated, yielding 2.80 g of a yellowish waxy solid, containing two products and some remaining starting material as shown by thin layer chromatographic analysis. The crude mixture was purified by medium pressure chromatography using a silica column and gradient elution, with stepwise increasing concentration of ethyl acetate in petroleum ether, resulting in 1.07 g of the title compound as a white solid.

¹H NMR (300 MHz, CDCl₃): δ 0.88 (t, J=7.2 Hz, 12 H), 1.26 (m, 96 H), 1.61 (m, 8 H), 2.30 (dd, J₁=7.7, J₂=3.2 Hz, 4 H), 2.32 (dd, J₁=7.4, J₂=3.7 Hz, 4H), 2.65-2.69 (m, 8 H), 4.14 (dd, J₁=11.8, J₂=3.0 Hz, 2H), 4.17 (dd, J₁=11.2, J₂=2.95 Hz, 2H), 4.27-4.34 (m, 4H), 5.23-5.29 (m, 2H), 5.77 (s, 2H). ¹³C NMR (300 MHz, CDCl₃): δ 14.114, 22.714, 24.890, 24.910, 28.556, 28.756, 29.116, 29.156, 29.303, 29.317, 29.390, 29.517, 29.530, 29.657, 29.671, 29.691, 29.731, 31.954, 34.057, 34.197, 62.018, 62.719, 68.775, 79.444, 170.858, 171.440, 172.849, 173.209.

20

g) Methylene bis[16-(15-(16-hydroxyhexadecanoyloxy-methoxycarbonyl)pentadecyloxy-carbonyl-ethylenecarbonyloxy)hexadecanoate

25

i) Methylene bis[16-(carboxyethylenecarbonyloxy)hexadecanoate]

Methylene bis(16-hydroxyhexadecanoate) (5.5g, 10mmol), succinic anhydride (2.2g, 22 mmol), and N,N-dimethyl-4-aminopyridine (2.5g, 20 mmol) were dissolved in N,N-dimethylformamide (100 ml) and heated to 50°C under stirring for 16 hours. N,N-dimethylformamide was evaporated in vacuo, the residue dissolved in chloroform/methanol (99:1) and filtered through a short silica column with the dissolving agent. The product was washed with petroleum ether, filtered, and dried

- 21 -

under vacuum. Yield: 6g white solid (78%). ^1H NMR (CDCl_3): δ 1.23-1.30 (m, 42H), 1.60-1.64 (m, 8H), 2.32-2.37 (t, 4H), 2.62-2.69 (m, 10H), 4.06-4.09 (t, 4H), 5.75 (s, 2H), 10-11 (m, 2H).

5

ii) Methylene bis[16-(15-(16-hydroxy-hexadecanoyloxymethoxycarbonyl)pentadecyloxy-carbonylethylenecarbonyloxy)hexadecanoate]

10 Methylene bis[16-(carboxyethylenecarbonyloxy)hexadecanoate] from (a) above (380mg, 0.5mmol), methylene bis(16-hydroxyhexadecanoate) (600mg, 1.1mmol), N-ethyl-N'-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (200mg, 1.05mmol), and N,N-dimethyl-4-aminopyridine (50mg) were dissolved in 25 ml N,N-dimethylformamide and stirred at 45°C for 16 hours. N,N-dimethylformamide was evaporated in vacuo, the residue dissolved in chloroform and purified by column chromatography on Silica KG60 with
20 dichloromethane/methanol as eluant. Yield: 100 mg white solid; (10%). ^1H NMR (CDCl_3): δ 1.25 (m, 132H), 1.59-1.65 (m, 24H), 2.32-2.37 (t, 12H), 2.61 (s, 8H), 3.63 (t, 4H), 4.05-4.09 (t, 8H), 5.74 (s, 6H).

25 h) 2-Hexadecylmalonic acid dihexadecyl ester

1,8-Diazabicyclo[5.4.0]undec-7-ene (1,5-5) (4.567g, 30mmol) and 1-iodohexadecane (10.570g, 30mmol) were added to a solution of 2-hexadecylmalonic acid in N,N-dimethylformamide (75ml). After stirring at room
30 temperature for 22 hours the precipitate was filtered off and washed with N,N-dimethylformamide. The product was purified by flash chromatography on silica gel using chloroform/hexane as eluant. Yield: 52%. ^1H -NMR (300MHz, CDCl_3): δ 0.88 (t, 9H), 1.26 (m, 80H), 1.62 (m, 4H),
35 1.88 (m, 2H), 3.31 (t, 1H), 4.15 (m, 4H). FAB-MS: 799.7 (M+Na).

- 22 -

i) 2,2-Dihexadecylmalonic acid dihexadecyl ester

Sodium hydride (72mg, 3mmol) was added in portions to a cooled (ice/water) solution of 2-hexadecylmalonic acid dihexadecyl ester (2.33g, 3mmol) in tetrahydrofuran (45ml) under an argon atmosphere and the resulting mixture was stirred at ambient temperature for 30 minutes. The solvent was evaporated and the solid residue was redissolved in N,N-dimethylformamide (45ml). A solution of 1-iodohexadecane (1.06g, 3mmol) in N,N-dimethylformamide (20ml) was added and stirring was continued at room temperature for 24 hours. Water (75ml) was added to the reaction mixture after the pH had been adjusted to about 5 with 2N HCl. The mixture was extracted with chloroform (3x75ml) and the chloroform phase was washed with water (3x75ml). The organic phase was dried with magnesium sulphate and the solvent was removed at reduced pressure. The residue was purified by flash chromatography (silicagel 60, chloroform/hexane). Yield: 66%. ¹H-NMR (300MHz, CDCl₃): δ 0.88(t,12H), 1.26(m,108H), 1.60(m,4H), 1.85(m,4H), 4.09(t,4H). FAB-MS: 1024 (M+Na).

j) 3-Hexadecyloxycarbonyl-3-hydroxypentanedioic acid dihexadecyl ester

1,8-Diazabicyclo[5.4.0]undec-7-ene(1,5-5) (4.567g, 30mmol) was added at room temperature to a solution of citric acid (1.924g, 10mmol) in N,N-dimethylformamide (50ml). 1-Iodoheptadecane (10.570g, 30mmol) was added and the resulting mixture was stirred at 50°C for 21 hours. The precipitated product was filtered off and washed with N,N-dimethylformamide and the crude product was purified by flash chromatography on silica gel using chloroform/hexane for elution. Yield: 83%. ¹H-NMR (300MHz, CDCl₃): δ 0.88(t,9H), 1.26(m,78H), 1.60(m,6H), 2.84(q,4H), 4.07(t,4H), 4.13(s,1H), 4.20(t,2H). FAB-MS:

- 23 -

887.6 (M+Na) .

k) Methylene bis-16-[3-(2,3-bis-hexadecanoyloxypropoxy carbonyl)propanoyloxy] hexadecanoate

5

Succinic acid 2,3-bis-hexadecanoyloxypropyl monester (1.30 g, 1.94 mmol) and methylene bis-16-hydroxyhexadecanoate (515 mg, 0.925 mmol) and triphenylphosphine (510 mg, 1.94 mmol) were dissolved in tetrahydrofuran (25 ml) under a nitrogen atmosphere. To the resulting solution was added diethyldiazodicarboxylate (as a 38-40% solution in toluene; 930 μ l, 1.94 mmol). The resulting reaction mixture was stirred at ambient temperature for 24 hours then concentrated to dryness. The remaining solid was redissolved in toluene/methylene chloride (5:2) and charged on top of a short column of silica gel, gradient elution with increasing amount of ethyl acetate (from 0 to 40 %) in petroleum ether and evaporation of the solvents affording 1.43 g of the title compound (83%) as a white solid (Mp 47°C).

^1H NMR (300 MHz, CDCl_3): δ 0.88 (t, 12 H), 1.26 (m, 140 H), 1.62 (m, 16H), 2.28-2.38 (m, 12H), 2.61-2.65 (m, 8H), 4.08 (t, $J=7$ Hz, 4H), 4.14 (dd, $J_1=12$ Hz, $J_2=6$ Hz, 2H), 4.18 (dd, $J_1=12$ Hz, $J_2=6$ Hz, 2H), 4.29 (dd, $J_1=12$ Hz, $J_2=4.5$ Hz, 2 H) 4.32 (dd, $J_1=12$, $J_2=4.5$ Hz, 2H) 5.23-5.32 (m, 2H), 5.74 (s, 2H). ^{13}C NMR (300 MHz, CDCl_3): δ 14.12, 22.71, 24.64, 24.87, 24.90, 25.91, 28.62, 28.94, 29.00, 29.04, 29.10, 29.14, 29.25, 29.30, 29.38, 29.48, 29.50, 29.52, 29.56, 29.62, 29.64, 29.68, 29.72, 31.95, 34.00, 34.06, 34.20, 62.04, 62.58, 65.00, 68.79, 76.63, 77.05, 77.47, 79.05, 171.83, 172.11, 172.48, 172.87, 173.24. PD-MS: 1881 (M-H+23(Na)), 1603 (M-H+23(Na)-255), 906 (odd el. fragm. with H), 651 (even el fragm.). MALDI-MS: 1897 (M+39(K)), 1881 (M+23(Na)).

l) Ethylidene bis(16-acetoxylhexadecanoate)

- 24 -

i) Ethylidene bis(16-hydroxyhexadecanoate)

1,8-Diazabicyclo [5.4.0.]undec-7-ene (1,5-5) (2.74 g, 0.018 mol) was added to 16-hydroxyhexadecanoic acid
5 (4.90 g, 0.018 mol) in dimethylformamide (150 ml). After 5 minutes with stirring ethylidene iodide (2.54 g, 0.009 mol) was added and the mixture was left with stirring at 40°C for 3 days. The reaction mixture was cooled to 20°C and when precipitation was complete (2
10 hours) the precipitate was isolated by filtration. The product was treated with activated carbon and recrystallised twice from dichloromethane to give 1.03 g (20%) of the title compound. Differential scanning calorimetry indicated that onset melting temperature was
15 88.93°C. ¹H NMR (200 MHz, CDCl₃): δ 1.25 (s, 44H, CH₂), 1.45 (d, 3H, CH₃CH), 1.56 (m, 8H, CH₂), 2.30 (t, 4H, CH₂CO), 3.63 (t, 4H, 2 x CH₂O), 6.86 (q, 1H, CHCH₃). ¹³C NMR (50 MHz, CDCl₃): δ 20.86, 25.91, 26.98, 30.22, 30.44, 30.67, 30.84, 34.00, 35.30, 64.00, 89.00, 171.77
20 (C=O).

ii) Ethylidene bis-(16-acetoxylhexadecanoate)

Ethylidene bis-16-hydroxyhexadecanoate (911 mg, 25 1.74 mmol) and N,N-dimethyl-4-aminopyridine (32 mg) were dissolved in tetrahydrofuran (25 ml), whereafter triethylamine (0.6 ml) and acetic anhydride (1 ml) were added successively via a syringe under a nitrogen atmosphere. After stirring the reaction mixture for two
30 hours it was diluted with methylene chloride and washed successively with hydrochloric acid (1%), saturated aqueous sodium bicarbonate and water. The organic phase was dried, the solvent was evaporated and the solid residue was dried under vacuum for a period of 24 hours
35 yielding 1.07g (94%) of the title compound as a white solid (Mp 58°C).

¹H NMR (300 MHz, CDCl₃): δ 1.26 (m, 44 H), 1.46 (d,

- 25 -

J=5.5 Hz, 3H), 1.56-1.67(m, 8H), 2.04 (s, 6H), 2.27-2.33(m, 4H), 4.05 (t, J=6.7 Hz, 4H), 6.86 (q, J=5.5 Hz, 1H). ¹³CNMR (300 MHz, CDCl₃): δ 19.58, 21.01, 24.68, 25.93, 28.64, 29.02, 29.26, 29.28, 29.47, 29.54, 29.59, 29.61, 29.65, 34.13, 64.66, 76.62, 77.04, 77.47, 88.31, 171.19, 171.73.

m) Ethylidene bis-(16-hexadecanoyloxyhexadecanoate)

10 Ethylidene bis-16-hydroxyhexadecanoate (570 mg, 1mmol) was dissolved in 25 ml dichloromethane. Triethylamine (276 μl, 2 mmol) was added with stirring. Palmitoyl chloride (600 μl, 2 mmol) dissolved in dichloromethane (25 ml) was added dropwise with
15 stirring. The reaction mixture was stirred overnight, whereafter the solvent was evaporated in vacuo; the residue was dissolved in dichloromethane and chromatographed on a glass column packed with silica. The produced was eluted with hexane/dichloromethane
20 (1:2). Yield: 900 mg (85%).
¹H-NMR (300 MHz, CDCl₃): δ = 0.91-0.85(6H, m), 1.37-1.23 (91H, m), 1.47-1.45(3H, d), 1.67-1.55 (12H, m), 2.33-2.26(8H, m), 4.08-4.03(4H, t), 6.89-6.84(1H, t).

25 n) Methylene bis-(16-acetoxyhexadecanoate)

Methylene bis-16 hydroxyhexadecanoate (2.0 g, 3.47 mmol) and N,N-dimethyl-4-aminopyridine (50 mg) were dissolved in tetrahydrofuran (25 ml), whereafter
30 triethylamine (1.2 ml) and acetic anhydride (2 ml) were added successively via a syringe under a nitrogen atmosphere. After stirring the reaction mixture for two hours it was diluted with methylene chloride and washed successively with hydrochloric acid (1%), saturated
35 aqueous sodium bicarbonate and water. The organic phase was dried, the solvent was evaporated and the solid residue was dried under vacuum for a period of 24 hours

- 26 -

yielding 2.06 g (90%) of the title compound as a white solid.

¹H NMR (300 MHz, CDCl₃) δ 1.26 (m, 44H), 1.56-1.68 (m, 8H), 2.04 (s, 6H), 2.35 (t, J=7.5 Hz, 4H), 4.05 (t, J=6.7 Hz, 4H), 5.74 (s, 2H). ¹³C NMR (300 MHz, CDCl₃): δ 21.00, 24.64, 25.94, 28.64, 29.03, 29.24, 29.28, 29.46, 29.54, 29.59, 29.60, 29.65, 29.66, 34.00, 64.66, 76.62, 77.04, 77.47, 79.06, 171.18, 172.49.

10 o) Methylene bis-16-[3-(2,3-bis-hexadecanoyloxy propoxycarbonyl)propanoyloxy]dodecanoate

Succinic acid 2,3-bis-hexadecanoyloxypropyl monoester (324 mg, 0.485 mmol) and methylene bis-12-hydroxydodecanoate (135 mg, 0.304 mmol) and triphenylphosphine (127 mg, 0.485 mmol) were dissolved in tetrahydrofuran (5 ml) under a nitrogen atmosphere. To the resulting solution was added diethyldiazodicarboxylate as a 38-40% solution in toluene (233 μl, 0.485 mmol). The resulting reaction mixture was stirred at ambient temperature for 24 hours then concentrated to dryness. The remaining solid was redissolved in methylene/petroleum ether (4:1) and charged on top of a short column of silica gel, gradient elution with increasing amount of ethyl acetate (from 0 to 40%) in petroleum ether and evaporation of the solvents affording 283 mg (67%) of the title compound as a white solid.

¹H NMR (300 MHz, CDCl₃): δ 0.88 (t, 12H), 1.26 (m, 124H), 1.55-1.67 (m, 16H), 2.27-2.38 (m, 12H), 2.61-2.65 (m, 8H), 4.08 (t, J=7Hz, 4H), 4.14 (dd, J₁=12 Hz, J₂=6 Hz, 2H), 4.18 (dd, J₁=12 Hz, J₂=6Hz, 2H), 4.29 (dd, J₁=12 Hz, J₂=4.5 Hz, 2H), 4.32 (dd J₁=12 Hz, J₂=4.5 Hz, 2H), 5.23-5.32 (m, 2H), 5.74 (s, 2H). ¹³C NMR (300 MHz, CDCl₃): δ 14.13, 22.71, 24.62, 24.88, 24.90, 25.90, 28.60, 28.93, 29.00, 29.03, 29.10, 29.14, 29.23, 29.26, 29.30, 29.38, 29.42, 29.51, 29.68, 29.72, 31.95, 33.98, 34.06, 34.20,

- 27 -

62.04, 62.57, 64.98, 68.78, 76.63, 77.06, 77.48, 79.06,
171.84, 172.11, 172.46, 172.87, 173.24.

p) PEG 10000 methyl ether 16-hexadecanoylhexadecanoate

5

PEG 10000 methyl ether (7.500g, 0.75 mmol) was dissolved in toluene (140 ml) and pyridine (0.107g, 1.35 mmol) was added. The solution was heated to 60°C and 16-hexadecanoyloxyhexadecanoyl chloride (0.595g, 1.12 mmol) dissolved in toluene (10 ml) was added dropwise. The mixture was heated to reflux and after stirring under reflux for 3 days the mixture was cooled to room temperature and precipitated into hexane. After filtering, the precipitate was washed with hexane and dried. Flash chromatography on a silica column, eluting with 5% methanol in chloroform, gave 5.39g (68%) of the title compound. ¹H NMR (300 MHz, CDCl₃): δ (0.84 (t, CH₃), 1.21 (s (br), CH₂), 1.55-1.60 (m, CH₂), 2.20-2.35 (m, CH₂CO), 3.34 (s, CH₃O), 3.61 (s, OCH₂CH₂O), 4.01 (t, COOCH₂CH₂O), 4.18 (t, COOCH₂CH₂O). ¹³C NMR (75 MHz, CDCl₂): δ 13.94, 22.48, 24.70, 24.82, 25.73, 28.94, 29.05, 29.14, 29.26, 29.33, 29.39, 29.45, 31.71, 34.00, 58.84, 63.14, 68.99, 69.36, 69.86, 69.97, 70.01, 70.36, 70.74, 70.82, 70.86, 71.72, 77.10, 173.62, 173.80.

25

EXAMPLE 2 - PREPARATION OF GAS-FILLED MICROPARTICLES

General procedure

30

A solution of the wall-forming substance in toluene was prepared. This solution was added to three times its volume of water containing a surfactant and mixed with a high speed rotor-stator mixer (8000-25000 rpm for 60-240 seconds). The resulting emulsion was frozen using a methanol/dry ice bath and lyophilised, resulting in a white powder.

35

- 28 -

Example	Wall material	% wall material in toluene	Surfactant	surfactant (% in water)
2a	Stearic acid	10	Gum arabic	3
2b	Product Ex 1a	5	HSA	5
2c	Product Ex 1b	5	HSA	5
2d	Product Ex 1c	5	HSA	5
2e	Product Ex 1e	5	HSA	5
2f	Product Ex 1f	5	HSA	5
2g	Product Ex 1h	5	HSA	5
2h	Product Ex 1i	5	HSA	5
2i	Product Ex 1j	5	HSA	5
2j	Product Ex 1j	5	Ex. 1p	1
2k	Product Ex 1j	5 ¹⁾	HSA	5
2l	Product Ex 1j	5 ²⁾	HSA	5
2m	Product Ex 1k	5	HSA	5
2n	Product Ex 1l	5	HSA	5
2o	Product Ex 1m	5	HSA	5
2p	Product Ex 1n	5	HSA	5
2q	1-Octadecanol	5	HSA	5
2r	Trimethylol-ethane	5	HSA	5
2s	Cholesteryl palmitate	5	HSA	5

SUBSTITUTE SHEET (RULE 26)

- 29 -

2t	α -Palmitin	5	HSA	5
2u	α, β - Dipalmitin	5	HSA	5
2v	Tripalmitin	5	HSA	5
2w	2- Hexadecanone	5	HSA	5
2x	Stearyl stearate	5	HSA	5
2y	Nonadecanoic acid N- methylester	5	HSA	5

1) Xylene as solvent

2) Camphene as solvent

HSA = human serum albumin

EXAMPLE 3 - ACOUSTIC CHARACTERISATION (IN VITRO)

The lyophilised samples from Example 2 were re-dispersed in MilliQ water or 0.9% sodium chloride solution by shaking on a laboratory shaker for an appropriate time. The particles were visually inspected using a light microscope.

The acoustic effect of the suspensions was obtained by measuring ultrasonic transmission through suspensions of different concentrations in an aqueous matrix, using either a 3.5 MHz broadband transducer in a pulse-reflection technique or two transducers with centre frequencies 3.5 and 5 MHz, covering a range from 1.5 to 8 MHz. The aqueous solvent system was used as reference, and measurements were performed by stepwise dilution of the starting suspension with the carrier liquid. Measurements were done until the signal was

- 30 -

an acoustic attenuation of less than 0.1 db/cm.

	Example	Particles from Ex. 2 dispersed in (solvent)	Particle size (μm)	Result
5	3a	2a (H_2O)	2-10	Contrast
	3b	2b (H_2O)	2-10	Strong Contrast
	3c	2c (0.9% NaCl)	3-5	Strong Contrast
	3d	2d (0.9% NaCl)	3-5	Strong Contrast
	3e	2e (H_2O)	2-10	Strong Contrast
	3f	2f (0.9% NaCl)	1-5	Strong Contrast
10	3g	2g (0.9% NaCl)	2-10	Strong Contrast
	3h	2h (0.9% NaCl)	2-10	Strong Contrast
	3i	2i (0.9% NaCl)	2-10	Strong Contrast
	3j	2j (0.9% NaCl)	1-5	Contrast
	3k	2k (0.9% NaCl)	1-5	Strong Contrast
	3l	2l (0.9% NaCl)	2-5	Contrast
15	3m	2m (0.9% NaCl)	2-10	Strong Contrast
	3n	2n (0.9% NaCl)	1-10	Contrast
	3o	2o (0.9% NaCl)	2-10	Strong Contrast
	3p	2p (0.9% NaCl)	1-5	Contrast
	3q	2q (0.9% NaCl)	1-10	Contrast
	3r	2r (0.9% NaCl)	1-5	Contrast
20	3s	2s (0.9% NaCl)	2-10	Strong Contrast
	3t	2t (0.9% NaCl)	2-15	Contrast

- 31 -

3u	2u (0.9% NaCl)	2-10	Strong Contrast
3v	2v (0.9% NaCl)	2-10	Strong Contrast
3w	2w (0.9% NaCl)	2-10	Contrast
3x	2x (0.9% NaCl)	2-10	Strong Contrast
5 3y	2y (0.9% NaCl)	1-10	Strong Contrast

EXAMPLE 4 - ACOUSTIC CHARACTERISATION (IN VIVO)

10 General procedure

Dry microparticle powders prepared as in Example 2 were redispersed in a sterile 0.9% (wt/wt) sodium chloride (aq) solution by shaking on a laboratory shaker
15 for 12-16 hours.

The dispersions were injected in ear veins of chinchilla rabbits, and their contrast effect was measured using a Doppler technique in which an
20 ultrasound probe was placed directly on a carotid artery and the inferior caval vein. Signal height in Doppler units and duration in seconds were recorded. The obtained signal heights were significant, indicating a strong in vivo ultrasound contrast effect for the
25 dispersions. Long signal duration confirmed good in vivo stability.

30

- 32 -

Example 4	Particles from Example	Dry Matter [mg/ml]	Dose [mg/kg]	Artery		Vein	
				Peak (DU)	Duration (s)	Peak (DU)	Duration (s)
a	2b	10.6	1.8	2.6	14.7	2.7	159
b	2c	9.9	1.7	7.5	156	8.0	324
c	2d	9.9	1.7	6.0	96	5.5	252

5

- 33 -

Claims:

1. A microparticulate contrast agent comprising gas or
a gas precursor encapsulated by a non-polymeric and non-
5 polymerisable wall-forming material.
2. A contrast agent as claimed in claim 1 wherein the
wall-forming material is selected from fatty acids and
esters thereof, fatty alcohols and esters thereof, fatty
10 amines and amides thereof, lipophilic aldehydes and
ketones, lipophilic derivatives of sugars, cholic acids
and derivatives thereof, cholesterol and derivatives
thereof, aliphatic and aromatic hydrocarbons,
hydrophobically modified X-ray contrast agents and
15 biocompatible fat-soluble antioxidants.
3. A contrast agent as claimed in claim 1 which
contains one or more methylene diester units of Formula
- 20
$$\{(O)_m-CO-O-C(R^1R^2)-O-CO-(O)_n\} \quad (I)$$
- where R^1 and R^2 each represent a hydrogen atom or a
carbon-attached monovalent organic group or R^1 and R^2
together form a carbon-attached divalent organic group
25 and m and n are each selected from zero and 1.
4. A contrast agent as claimed in claim 3 wherein the
wall-forming material contains one or more methylene
diester units of formula (I) in which R^1 and R^2 are
30 selected from hydrogen atoms and methyl groups and n and
 m are both zero.
5. A contrast agent as claimed in any of the preceding
claims further comprising a polymeric emulsifier.
- 35 6. A contrast agent as claimed in claim 5 wherein the
polymeric emulsifier is selected from polyvinyl alcohol,

- 34 -

- proteins, polysaccharides, block copolymers consisting of alternating hydrophilic and hydrophobic blocks, polyoxyethylated derivatives of partial fatty acid esters of hexahydric alcohols, polyvinylpyrrolidones, fatty alcohol ethers of polyoxyethylene alcohols, polyethylene glycol esters of fatty acids, polyethylene glycol-sorbitan-beeswax surfactants and perfluorinated derivatives of any of the above polyethylene glycol-containing compounds.
7. A contrast agent as claimed in claim 6 wherein the polymeric emulsifier is human serum albumin.
8. Use of a contrast agent as claimed in any of the preceding claims in diagnostic imaging.
9. Use of a contrast agent as claimed in any of claims 1 to 7 in diagnostic ultrasound imaging.
10. Use of a contrast agent as claimed in any of claims 1 to 7 in magnetic resonance imaging.
11. A method of generating enhanced images of a human or non-human animal body which comprises administering to said body a contrast agent as claimed in any of claims 1 to 7 and generating an ultrasound or magnetic resonance image of at least a part of said body.
12. A process for the preparation of a contrast agent as claimed in claim 1 which comprises (i) generating an emulsion comprising hydrophilic and hydrophobic phases wherein wall-forming material is preferentially solubilised in the dispersed phase or about the interfaces between the phases and (ii) isolating the desired contrast agent from said emulsifier.

- 35 -

13. A process as claimed in claim 12 effected in the presence of at least one emulsifier.

14. A process as claimed in claim 13 wherein said
5 emulsifier is a polymeric emulsifier.

INTERNATIONAL SEARCH REPORT

Internal Application No. PCT/GB 95/00316

A. CLASSIFICATION OF SUBJECT MATTER

A 61 K 49/00, A 61 K 49/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A 61 K, A 61 B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT*

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, F	WO, A, 94/09 703 (DREXEL UNIVERSITY) 11 May 1994 (11.05.94), abstract; claims.	1, 2, 12
X	EP, A, 0 365 467 (SCHERING AKTIENGESSELLSCHAFT) 25 April 1990 (25.04.90), abstract; claims.	1
A	WO, A, 93/17 718 (NYCOMED IMAGING A/S) 16 September 1993 (16.09.93), abstract; claims.	1, 3
A	WO, A, 94/01 140 (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH)	1, 2, 12



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search
25 April 1995

Date of mailing of the international search report

17.05.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Mme Dagmar FRANK

INTERNATIONAL SEARCH REPORT

-2-

International Application No

PCT/GB 95/00316

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	<p>20 January 1994 (20.01.94), abstract; claims. -----</p>	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 95/00316

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 8-11
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 8-11 are disclosed by Article 17(2)(a)i) with Rule 39.1(iv) PCT these claims have been searched.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6A(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANHANG

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

ANNEX

to the International Search
Report to the International Patent
Application No.

ANNEXE

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/GB 95/00316 SAE 104185

In diesem Anhang sind die Mitglieder
der Patentfamilien der in obenge-
nannten internationalen Recherchenbericht
angeführten Patentedokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visé ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

In Recherchenbericht angeführtes Patentedokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
WD A1 9409703	11-05-94	AU A1 54573/94	24-05-94
		US A 5352436	04-10-94
EP A2 365467	25-04-90	AT E 93145	15-09-93
		AU A1 33900/89	12-04-90
		AU B2 627456	27-08-92
		CN A 1044225	01-08-90
		DE A1 3834705	12-04-90
		DE C0 58905318	23-09-93
		DK A0 997/89	01-03-89
		DK A 997/89	08-04-90
		EP A3 365467	06-06-90
		EP B1 365467	18-08-93
		ES T3 2059814	16-11-94
		FI A0 890749	16-02-89
		FI A 890749	08-04-90
		IL A0 90028	15-12-89
		IL A1 90028	31-01-93
		JP A2 2145528	05-06-90
		NO A0 890609	13-02-89
		NO A 890609	09-04-90
		NZ A 228781	26-03-92
		PT A 91523	30-04-90
		US A 5141738	25-08-92
		ZA A 8902885	28-12-90
		AT E 36959	15-09-88
		AU A1 26806/84	18-10-84
		AU B2 569072	21-01-88
		CA A1 1232837	16-02-88
		DE A1 3313947	18-10-84
		DE C0 3473829	13-10-88
		DK A0 790/84	20-02-84
		DK A 790/84	16-10-84
		DK B 165622	28-12-92
		DK C 165622	01-06-93
		EP A2 123235	31-10-84
		EP A3 123235	20-11-86
		EP B1 123235	07-09-88
		FI A0 841463	12-04-84
		FI A 841463	16-10-84
		FI B 81265	29-06-90
		FI C 81265	10-10-90
		IE B 57197	03-06-92
		JP A2 59205329	20-11-84
		JP B4 4017164	25-03-92
		NO A 841490	16-10-84
		NO B 163669	26-03-90
		NO C 163669	04-07-90
		NZ A 207854	08-01-88
		ZA A 8402800	28-11-84
		AT E 36958	15-09-88
		AU A1 26805/84	18-10-84
		AU B2 566928	05-11-87
		CA A1 1239092	12-07-88
		DE A1 3313946	18-10-84
		DE C0 3473828	13-10-88
		DK A0 789/84	20-02-84
		DK A 789/84	16-10-84
		DK B 165171	19-10-92
		DK C 165171	01-03-93
		EP A2 122624	24-10-84
		EP A3 122624	20-11-86
		EP B1 122624	07-09-88
		FI A0 841462	12-04-84

FI A	841462	16-10-84
FI B	81264	29-06-90
FI C	81264	10-10-90
IE B	57272	01-07-92
JP A2	59205328	20-11-84
JP B4	4025934	06-05-92
NO A	841489	16-10-84
NO B	161356	02-05-89
NO C	161356	09-08-89
NZ A	207853	08-01-88
ZA A	8402801	28-11-84

WD A1 9317718 16-09-93

AU A1	36424/93	05-10-93
CN A	1084410	30-03-94
EP A1	627942	14-12-94
FI A	944077	05-09-94
FI A0	944077	05-09-94
GB A0	9204920	22-04-92
HU A0	9402565	28-11-94
IL A0	104963	08-07-93
NO A	943272	05-09-94
NO A0	943272	05-09-94
AU A1	36423/93	05-10-93
CN A	1084524	30-03-94
EP A1	629214	21-12-94
FI A	944078	05-09-94
FI A0	944078	05-09-94
GB A0	9204918	22-04-92
HU A0	9402563	28-11-94
IL A0	104962	08-07-93
NO A	943273	05-09-94
NO A0	943273	05-09-94
WD A1	9318070	16-09-93

WD A1 9401140 20-01-94

AU A1 45630/93 31-01-94